# Overactive Bladder Medications

# **Drug Class: Antichoinergics**

### **Medications:**

Toviaz (fesoterodine)

Trospium (sanctura)

Vesicare (solifenacin)

Enablex (darifenacin)

Detrol/Detrol LA(tolterodine)

Ditropan (oxybutynin)

## **Common Side Effects:**

dry mouth constipation headaches urinary retention

## **Serious Side Effects:**

confusion/dementia

**Drug Class: Beta Agonists** 

### **Medications:**

Myrbetriq (Mirabegron) Gemtesa (Vibegron)

#### **Common Side Effects:**

elevated blood pressure diarrhea urinary retention

If successful, medication can take up to 4 weeks to show improvement.

Medication may be started simutaneously with lifestyle adjustments.

## **Medication Adjustments**

Adjustments can be made to the

- •dose (mg)
- •frequency (daily, twice a day, or at bedtime)
- medication (switching medications)

## **Insurance Coverage**

Insurance does affect which medications are covered and how much they will cost.

This varies by situation and insurance policy. This can change every year depedning on your policy. If you have changes to your policy and this affects your medication or treatment, please notify us.

Nurse Navigator: Ashley Phone: 573-381-5207 Fax: 573-335-2007

Fax: 573-335-2007 Email: atilley@capeurology.net

